

Metal Sulfides and their Relation to Atmospheric Sulfur on Venus

S.T. PORT, A.C. BRISCOE, V.F. CHEVRIER

ARKANSAS CENTER FOR SPACE AND PLANETARY SCIENCES


THIS STUDY WAS SUPPORTED BY NASA SOLAR SYSTEM WORKINGS GRANT #NNX15AL57G



Introduction

2

- ▶ Sulfur is an important constituent in the atmosphere
 - ▶ SO_2
 - ▶ COS
 - ▶ H_2SO_4
- ▶ More abundant in atmosphere than on Earth
- ▶ Expect a complex Sulfur Cycle on Venus
- ▶ Little understanding of the surface composition
- ▶ Sources and sinks of sulfur?



Fegley, B., et al. (1995)

Objective

3

- ▶ **Determine possible sources and sinks for sulfur:**
 - ▶ Venusian temperature and pressure
 - ▶ CO_2 , SO_2 , and COS

Mineralogy

4

▶ Galena (PbS)

- ▶ **SO₂ can be released via the oxidation** Abdel-Rehim, A.M., 2006
- ▶ **Most common lead mineral on Earth** Nowak, P. et al., 2009
- ▶ **On list of metal frost candidates** Schaefer, L., et al., 2004

▶ Pyrrhotite (Fe₇S₈)

- ▶ **Speculated to be one of the most abundant sulfur minerals on Venus** Fegley, B., et al., 1992
- ▶ **Decomposition can release COS** Fegley, B., et al., 1995
- ▶ **On list of metal frost candidates** Fegley, B., et al., 1992

▶ Metacinnabar (HgS)

- ▶ **Stable form of cinnabar at high temperatures** Ballirano, P., et al., 2013
- ▶ **Temperature sensitive** Ballirano, P., et al., 2013
- ▶ **Found near volcanic activity** Rytuba J.J. et al., 1992

Methods

5

- ▶ One gram of each mineral
- ▶ Two Scenarios:
 - ▶ 1. Oven
 - ▶ Lindberg Tube Oven
 - ▶ Temperature
 - ▶ 460°C (avg. lowland altitude)
 - ▶ 425°C (slightly above frost line)
 - ▶ 380°C (11 km)
 - ▶ Gases
 - ▶ CO₂
 - ▶ CO₂ 100ppm SO₂
 - ▶ CO₂ 100ppm COS



Methods

6

▶ 2. Chamber

▶ UArk Cassiopeia Chamber

▶ Temperature/Pressure

▶ 460°C/95 bar

▶ 425°C/75 bar

▶ 380°C/45 bar

▶ Gases

▶ CO₂

▶ CO₂ 100ppm SO₂

▶ CO₂ 100ppm COS

▶ All experiments lasted 24 hours

▶ All samples were analyzed with the PANalytical X'Pert MRD



Results

7



- Pyrrhotite: Untreated (left), 380°C in CO₂, 425°C in CO₂, 460°C in CO₂ (right)

Pyrrhotite CO₂ Oven v. Chamber

8

	460°C/1 bar (lowlands)	425°C/1 bar (frost line)	380°C/1 bar (highlands)
Oven	Hematite (Fe₂O₃) Mikasaite (Fe₂(SO₄)₃)	Magnetite (Fe₃O₄) Pyrrhotite (Fe₇S₈)	Pyrrhotite (Fe₇S₈) Troilite (FeS)
	460°C/95 bar	425°C/75 bar	380°C/45 bar
Chamber	Pyrrhotite Troilite	-----	Pyrrhotite Troilite

Pyrrhotite SO_2 v. COS (Oven)

9

	460°C/1 bar (lowlands)	425°C/1 bar (frost line)	380°C/1 bar (highlands)
CO_2/SO_2	Pyrrhotite Troilite Hematite	----	Pyrite (FeS_2) Pyrrhotite Hematite Troilite Magnetite
	460°C/1 bar	425°C/1 bar	380°C/1 bar
CO_2/COS	Hematite Mikasaite	Hematite Maghemite Mikasaite	Pyrrhotite Pyrite Hematite

Galena CO₂ Oven v. Chamber

10

	460°C/1 bar (lowlands)	425°C/1 bar (frost line)	380°C/1 bar (highlands)
Oven	Galena (PbS) Anglesite (Pb(SO ₄)) Lanarkite (Pb ₂ (SO ₄)O)	Galena (PbS) Anglesite (Pb(SO ₄)) Lanarkite (Pb ₂ (SO ₄)O)	Galena (PbS) Anglesite (Pb(SO ₄))
	460°C/95 bar	425°C/75 bar	380°C/45 bar
Chamber	Galena	-----	Galena PbO (Litharge)


Galena SO_2 v. COS (Oven)

11

	460°C/1 bar (lowlands)	425°C/1 bar (frost line)	380°C/1 bar (highlands)
CO_2/SO_2	Galena Anglesite Lanarkite	Galena Anglesite	Galena Anglesite
	460°C/1 bar	425°C/1 bar	380°C/1 bar
CO_2/COS	Galena Anglesite	Galena Anglesite	Galena Anglesite

Metacinnabar

12



	380°C/1 bar (highlands)
CO ₂ /SO ₂	Metacinnabar
	380°C/1 bar
CO ₂ /COS	Cinnabar

Pyrrhotite

13

- ▶ **Pyrrhotite** → **Magnetite** → **Maghemite** → **Hematite**

Fegley, B., et al., 1995

- ▶ **Troilite**: Vaporization of S increases the ratio of Fe to S

- ▶ Quicker oxidization in mixed gas experiments

- ▶ Pyrite formation in the low temperature, mixed gas experiments


- ▶ Product of oxidation

- ▶ $3\text{Fe}_7\text{S}_8 + 28\text{CO}_2 \leftrightarrow 7\text{Fe}_3\text{O}_4 + 12\text{S}_2 + 28\text{CO}$



Fegley, B., et al., 1995

- ▶ Unable to verify




Mycroft, J. R., et al. (1994)

Galena

14

- ▶ Formation of Anglesite:



- ▶ Formation of Lanarkite:



- ▶ Formation of Lead Oxide (Litharge):



- ▶ SO_2 produced in all equations

- ▶ Currently unable to verify

Metacinnabar

15

- ▶ **Instability in all CO₂ experiments in the oven**
- ▶ **Cinnabar is a low T/P version of metacinnabar**
- ▶ **Heating and cooling of metacinnabar can form cinnabar**
Ballirano, P., et al., 2013
- ▶ **Stability in CO₂ in the chamber at lowland and highland conditions**

Future Work

16

- ▶ **Gas Chromatograph**
- ▶ **Gas Mixture Experiments in the Chamber**
- ▶ **In situ Studies with RAMAN**
- ▶ **Longer Experiments (48-72h)**

Conclusion

17

- ▶ **Pyrrhotite**
 - ▶ **Unstable in oven**
 - ▶ **Stable in chamber**
 - ▶ **More rapid oxidation in mixed gases**

Conclusion

17

- ▶ **Pyrrhotite**
 - ▶ Unstable in oven
 - ▶ Stable in chamber
 - ▶ More rapid oxidation in mixed gases
- ▶ **Galena**
 - ▶ Minor instability in oven
 - ▶ Better stability in chamber
 - ▶ Mixed gases had no effect

Conclusion

17

- ▶ **Pyrrhotite**
 - ▶ Unstable in oven
 - ▶ Stable in chamber
 - ▶ More rapid oxidation in mixed gases
- ▶ **Galena**
 - ▶ Minor instability in oven
 - ▶ Better stability in chamber
 - ▶ Mixed gases had no effect
- ▶ **Metacinnabar**
 - ▶ Unstable in high temperatures in oven
 - ▶ May show better stability in chamber

Conclusion

17

- ▶ **Pyrrhotite**
 - ▶ Unstable in oven
 - ▶ Stable in chamber
 - ▶ More rapid oxidation in mixed gases
- ▶ **Galena**
 - ▶ Minor instability in oven
 - ▶ Better stability in chamber
 - ▶ Mixed gases had no effect
- ▶ **Metacinnabar**
 - ▶ Unstable in high temperatures in oven
 - ▶ May show better stability in chamber
- ▶ **Mixed gas experiments need to be completed in the chamber**
- ▶ **Currently cannot determine what gases are released during reactions**
 - ▶ **Source/Sink?**